

Superfield formulation of $\mathcal{N} = 4$ Super Yang-Mills theory in plane wave background*

R.R. Metsaev[†]

*Department of Theoretical Physics, P.N. Lebedev Physical Institute,
Leninsky prospect 53, Moscow 119991, Russia*

Abstract

The superfield light cone gauge formulation of $\mathcal{N} = 4$ SYM in plane wave background is developed. We find a realization of superconformal symmetries in terms of a light cone superfield. An oscillator realization of superconformal symmetries is also obtained.

*Contribution to Proceedings of the 3rd International Sakharov conference on physics. June 24-29, Moscow, Russia.

[†]E-mail: metsaev@lpi.ru

1 Introduction

The new maximally supersymmetric solution of IIB supergravity with Ramond-Ramond flux [1] attracted recently considerable interest. On the one hand, the light cone gauge Green-Schwarz superstring action in this background is quadratic in both bosonic and fermionic superstring $2d$ fields, and therefore, this model can be explicitly quantized [2]. On the other hand, in [3] it was proposed that this superstring in plane wave background corresponds to a certain (large R-charge) sector of the $\mathcal{N} = 4$ SYM theory. Given that the plane wave superstring model can be quantized explicitly [2, 4] one can study the duality correspondence between string states and gauge theory operators at the string-mode level [3]. The new duality [3] turned out to be very fruitful and this renewed interest in various aspects of string/gauge theory correspondence.

One of the important aspects of the string/gauge theory correspondence is to find explicit and precise relation between string states and appropriate operators of gauge theory. For the case of plane wave duality solution to this problem was suggested in [3] by introducing the operators which are often referred to as BMN operators. These operators were extensively studied in the literature (see for instance [5, 6, 7] and references in [8]) and various improvements of the original BMN operators were suggested [9, 10, 11, 8, 12]. Unfortunately, the guiding principles which would allow one to fix the precise form of these operators are not clear so far¹. Recent interesting discussion of this theme may be found in [18]. One of the important inputs in establishing a precise correspondence might be the geometry of space-time in which fields of SYM are propagating. Usually, the Hamiltonian formulation of conformal field theory for SYM is considered in $R \times S^3$ background². The space-time symmetries of this background, however, are in sharp contradiction with the (super)symmetries of the plane wave superstring model. Therefore, taking into account that some (super)symmetries of plane wave SYM action are look the same as (super)symmetries of the type IIB superstring in plane wave Ramond-Ramond background it was suggested [22] that the SYM theory in plane wave background may be preferable in the context of the new duality.

Another argument in favor of the discussion of the plane wave SYM is that $4d$ plane wave background can be obtained via Penrose limit from $R \times S^3$ background. Given that the SYM in $R \times S^3$ is used for study of holography in the original context of AdS/CFT correspondence we expect that $4d$ plane wave background is most appropriate for the study of holographical issues of the plane wave duality. Thus, it seems natural to expect that in order to match the states of the plane wave superstring and operators of SYM the Penrose limit on the string theory side should be supplemented by the Penrose limit on the SYM side. Various alternative discussions of plane wave holography may be found in [23].

In [22] we developed the Hamiltonian light cone gauge formalism for the plane wave SYM in component form. In this paper we continue the study of the plane wave SYM and develop a superfield formulation of this theory. Superfield formulation turns out to be

¹It seems natural to expect that these operators should be derivable from the microscopic open-string approaches to the supersymmetric plane wave Dp branes. Discussion Dp branes in the framework of microscopic approach may be found in [13, 14, 15, 16, 17].

²Discussion of SYM theory in various curved backgrounds may be found in [19]. Plane wave SYM does not fall in the cases considered in [19]. Detailed study of the Hamiltonian formulation of SYM in $R \times S^3$ background may be found in [20]. Relevance of a curved background for SYM was discussed (in the original AdS/CFT context) in [21].

more compact, and we expect that this formalism will be helpful in future studies (similar to one in [24]). Independently of the above string/gauge theory duality motivation it seems interesting to study SYM theory in curved backgrounds and the present work may be also considered as an effort in this direction.

2 Action of plane wave $\mathcal{N} = 4$ SYM in light cone superspace

Component form of light cone gauge action of $\mathcal{N} = 4$ SYM is formulated in terms bosonic fields $A^I = (A^{\hat{i}}, \phi^M)$, $A^{\hat{i}}$ is spin one field and ϕ^M are six scalar fields, and 16-component fermionic field $\psi^{\oplus}$ which satisfies the constraint $\bar{\gamma}^- \psi^{\oplus} = 0$. Because on plane wave Ramond-Ramond superstring theory side $so(8)$ symmetries are broken to $so(4) \oplus so'(4)$ it is natural to reduce manifest SYM R-symmetries $so(6)$ to the R-symmetries $so(2) \oplus so'(4)$. Another technical advantages in reducing manifest SYM R-symmetries is related with the fact in this way one can formulate SYM theory in terms of unconstrained light cone superfield. As in the flat space-time the superfield formulation of $\mathcal{N} = 4$ SYM could be developed by using a certain superspace whose 16 real-valued fermionic coordinates related with the θ^α which form Glifford algebra

$$\{\theta^\alpha, \theta^\beta\} = \frac{1}{4}(\gamma^+)^{\alpha\beta}, \quad \bar{\gamma}^+ \theta = 0. \quad (2.1)$$

However the superfield based on such fermionic coordinates should satisfy complicated constraints [25]. To avoid such superfield description one needs to introduce Grassmann coordinates instead of Clifford coordinates θ^α [26, 27]. To this end we break manifest R-symmetries $so(6)$ to R-symmetries $so(2) \oplus so'(4)$ and introduce the fermionic ‘coordinates’ θ^Z and fermionic ‘momenta’ $\theta^{\bar{Z}}$ by relations

$$\theta^Z = \frac{1}{2}\gamma^{\bar{Z}}\bar{\gamma}^Z\theta, \quad \theta^{\bar{Z}} = \frac{1}{2}\gamma^Z\bar{\gamma}^{\bar{Z}}\theta, \quad \gamma^Z = \frac{1}{\sqrt{2}}(\gamma^3 + i\gamma^4), \quad \gamma^{\bar{Z}} = \gamma^{Z*}, \quad (2.2)$$

which satisfy the desired commutation relations

$$\{\theta^{Z\alpha}, \theta^{\bar{Z}\beta}\} = \frac{1}{8}(\gamma^+\gamma^{\bar{Z}}\gamma^Z)^{\alpha\beta}, \quad \{\theta^{Z\alpha}, \theta^{Z\beta}\} = 0, \quad \{\theta^{\bar{Z}\alpha}, \theta^{\bar{Z}\beta}\} = 0. \quad (2.3)$$

Accordingly the fields of SYM are also decomposed into multiplets of R-symmetry $so(2) \oplus so'(4)$ algebra. The six scalar fields ϕ^M are decomposed into four scalar fields $\phi^{i'}$ transforming in vector representation of the $so'(4)$ algebra and two complex fields $Z, \bar{Z}$ defined by

$$Z \equiv \frac{1}{\sqrt{2}}(\phi^3 + i\phi^4), \quad \bar{Z} \equiv \frac{1}{\sqrt{2}}(\phi^3 - i\phi^4), \quad (2.4)$$

while the fermionic field $\psi^{\oplus}$ is decomposed as follows

$$\psi^{\oplus} = \psi^{\oplus Z} + \psi^{\oplus \bar{Z}}, \quad \psi^{\oplus Z} \equiv \frac{1}{2}\gamma^{\bar{Z}}\bar{\gamma}^Z\psi^{\oplus}, \quad \psi^{\oplus \bar{Z}} \equiv \frac{1}{2}\gamma^Z\bar{\gamma}^{\bar{Z}}\psi^{\oplus}. \quad (2.5)$$

Light-cone gauge $\mathcal{N} = 4$ SYM in plane wave background can be formulated in light-cone superspace which is based on space-time coordinates $x^\pm, x^{\hat{i}}$ and Grassmann coor-

dinates θ^Z ³. In this light-cone superspace we introduce scalar superfield $\phi(x^\pm, x^{\hat{i}}, \theta^Z)$. Instead of position space it is convenient to use momentum space for all coordinates except the light-cone time x^+ . This implies using momenta $p^+, p^{\hat{i}}, \theta^Z$, instead of positions $x^-, x^{\hat{i}}, \theta^Z$ respectively. Thus we consider the scalar superfield $\phi(x^+, p^+, p^{\hat{i}}, \theta^Z)$ with the following expansion in powers of Grassmann momentum θ^Z

$$\begin{aligned}\phi(p, \theta^Z) &= p^+ \bar{Z} + i\theta^{\bar{Z}} \bar{\gamma}^{\bar{Z}} \psi^{\oplus \bar{Z}} - \frac{1}{2} \theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z} \hat{i}} \theta^{\bar{Z}} A^{\hat{i}} - \frac{1}{2} \theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z} i'} \theta^{\bar{Z}} \phi^{i'} \\ &- \frac{i}{6p^+} \theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z} \hat{i}} \theta^{\bar{Z}} \theta^{\bar{Z}} \bar{\gamma}^{\hat{i}} \psi^{\oplus Z} - \frac{1}{24p^+} (\theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z} \hat{i}} \theta^{\bar{Z}})^2 Z.\end{aligned}\quad (2.6)$$

The fact that the fields $A^I(p)$, $\psi^{\oplus}(p)$ are subject to the hermitian conjugation rules given in (4.51) can be expressed in terms of the superfield by the following relation

$$\phi(-p, \theta^{\bar{Z}}) = (p^+)^2 \int d^4 \theta^{\bar{Z} \dagger} e^{2\theta^{\bar{Z}} \bar{\gamma} - \theta^{\bar{Z} \dagger} / p^+} \phi(p, \theta^{\bar{Z}})^\dagger. \quad (2.7)$$

In terms of the superfield $\phi(p, \theta^{\bar{Z}})$ the action of $\mathcal{N} = 4$ SYM takes then the form

$$S = S_2 + S_3 + S_4, \quad (2.8)$$

where the free action S_2 and cubic action S_3 are fixed to be⁴

$$S_2 = \text{Tr} \int d^4 \theta^{\bar{Z}} dx^+ d^3 p p^+ \phi(-p, -\theta^{\bar{Z}}) (i\partial_{x^+} + P^-) \phi(p, \theta^{\bar{Z}}), \quad (2.9)$$

$$S_3 = \text{Tr} \frac{1}{3} \int d\Gamma_3 \mathbf{P}^{\hat{i}} \frac{\Theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z} \hat{i}} \Theta^{\bar{Z}}}{p_1^+ p_2^+ p_3^+} \prod_{a=1}^3 \phi(p_a, \theta_a^{\bar{Z}}). \quad (2.10)$$

The expression for the particle Hamiltonian P^- which enters the free action is given in (3.19), while the integration measure in cubic action is given by

$$d\Gamma_3 = \frac{1}{(2\pi)^{3/2}} \delta^{(3)} \left(\sum_{a=1}^3 p_a \right) \delta \left(\sum_{a=1}^3 \theta_a^{\bar{Z}} \right) \prod_{a=1}^3 d^3 p_a d^4 \theta_a^{\bar{Z}}. \quad (2.11)$$

The quantities $\mathbf{P}^{\hat{i}}$, $\Theta^{\bar{Z}}$ are defined by

$$\mathbf{P}^{\hat{i}} \equiv \frac{1}{3} \sum_{a=1}^3 p_a^{\hat{i}} (p_{a+1}^+ - p_{a+2}^+), \quad \Theta^{\bar{Z}} \equiv \frac{1}{3} \sum_{a=1}^3 \theta_a^{\bar{Z}} (p_{a+1}^+ - p_{a+2}^+), \quad (2.12)$$

so that they are manifestly symmetric under cyclical permutations of particle labels 1,2,3. Taking into account the momenta conservation laws we obtain the relations

$$\mathbf{P}^{\hat{i}} = \mathbf{P}_{12}^{\hat{i}} = \mathbf{P}_{23}^{\hat{i}} = \mathbf{P}_{31}^{\hat{i}}, \quad \Theta^{\bar{Z}} = \Theta_{12}^{\bar{Z}} = \Theta_{23}^{\bar{Z}} = \Theta_{31}^{\bar{Z}}, \quad (2.13)$$

³ $\mu = 0, 1, \dots, 9$ are $so(9, 1)$ vector indices; $I, J, K = 1, \dots, 8$ are $so(8)$ transverse indices; $\hat{i}, \hat{j}, \hat{k} = 1, 2$ are space-time symmetry $so(2)$ transverse indices; $M, N = 3, \dots, 8$ are R-symmetry $so(6)$ indices; $i', j' = 5, 6, 7, 8$ are R-symmetry $so'(4)$ indices; indices $\hat{I}, \hat{J} = 1, 2, 5, 6, 7, 8$ are collection of space-time $so(2)$ indices $\hat{i}, \hat{j}$ and R-symmetry $so'(4)$ indices i', j' . Coordinates in light-cone directions are defined by $x^\pm \equiv (x^9 \pm x^0)/\sqrt{2}$. Remaining transverse coordinates x^I are decomposed into $x^{\hat{i}}, x^{i'}, x^{Z, \bar{Z}}$ where $x^{Z, \bar{Z}} \equiv (x^3 \pm ix^4)/\sqrt{2}$. The scalar product of two $so(8)$ vectors is decomposed then as $X^I Y^I = X^{\hat{i}} Y^{\hat{i}} + X^{i'} Y^{i'} + X^Z Y^Z + X^{\bar{Z}} Y^{\bar{Z}}$.

⁴Because the expression for S_4 is not illuminating we do not present it here.

where

$$\mathbf{P}_{ab}^{\hat{i}} \equiv p_a^{\hat{i}} p_b^+ - p_b^{\hat{i}} p_a^+, \quad \Theta_{ab}^{\bar{Z}} \equiv \theta_a^{\bar{Z}} p_b^+ - \theta_b^{\bar{Z}} p_a^+. \quad (2.14)$$

The superfield form of the action (2.9),(2.10) was obtained by direct comparison with the action taken in terms of component fields (4.45),(4.48).

In the Hamiltonian approach the physical fields satisfy the canonical (anti)commutation relations

$$[A^I(p)^a, A^J(p')^{a'}]_{\text{equal } x^+} = \frac{1}{2p^+} \delta^{IJ} \delta^{(3)}(p + p') I^{aa'}, \quad (2.15)$$

$$\{\psi^{\oplus Z\alpha}(p)^a, \psi^{\oplus \bar{Z}\beta}(p')^{a'}\}_{\text{equal } x^+} = -\frac{1}{4} (\gamma^- \bar{\gamma}^{\bar{Z}} \gamma^Z)^{\alpha\beta} \delta^{(3)}(p + p') I^{aa'}, \quad (2.16)$$

where $I^{aa'}$ is a projector operator which we insert to respect the Lie algebra indices of the physical fields $A^I = A^{Ia} t_a$ and $\psi^{\oplus} = \psi^{\oplus a} t_a$. All that is required this operator should satisfy the relation $I^{ac} \text{Tr}(t_c t_b) = \delta_b^a$. Taking into account the expression for δ -function over anti-commuting variables (4.79) the above-given canonical commutation relations can be collected into the superfield form

$$[\phi(p, \theta^{\bar{Z}})^a, \phi(p', \theta^{\bar{Z}'})^{a'}]_{\text{equal } x^+} = \frac{1}{2p^+} \delta^{(3)}(p + p') \delta(\theta^{\bar{Z}} + \theta^{\bar{Z}'}) I^{aa'}. \quad (2.17)$$

3 Superfield realization of superconformal symmetries of $\mathcal{N} = 4$ plane wave SYM

In this section we discuss realization of global superconformal supersymmetries of $\mathcal{N} = 4$ SYM in plane wave background. As is well known these symmetries are generated by the $psu(2, 2|4)$ superalgebra. The realization of these symmetries on the component fields of $\mathcal{N} = 4$ SYM was obtained in [22]. Here we develop a superfield realization of these symmetries. To do that we use the framework of Noether charges. The Noether charges play an important role in analysis of the symmetries of dynamical systems. The choice of the light cone gauge spoils manifest global symmetries, and in order to demonstrate that these global invariances are still present, one needs to find the Noether charges that generate them.

The component form of the Noether charges for $\mathcal{N} = 4$ SYM was obtained in [22]. Superfield representation for the Noether charges can be derived by direct comparison with the Noether charges taken in the component form. Doing this we obtain the following representation for the Noether charges $G_{f.t.}$.

$$G_{f.t.} = \int d^4 \theta^{\bar{Z}} d^3 p p^+ \phi(-p, -\theta^{\bar{Z}}) G \phi(p, \theta^{\bar{Z}}), \quad (3.18)$$

where G denotes representation of generators of the $psu(2, 2|4)$ superalgebra in terms of differential operators acting on the superfield $\phi(p, \theta^{\bar{Z}})$. These differential operators are listed below.

Generators of the isometry symmetries:

$$P^+ = p^+, \quad P^- = -\frac{p^2}{2p^+} + \frac{f^2}{2} p^+ \partial_p^2, \quad (3.19)$$

$$T^{\hat{i}} = e^{-\text{if}x^+}(p^{\hat{i}} - \text{f}p^+ \partial_{p^{\hat{i}}}), \quad \bar{T}^{\hat{i}} = e^{\text{if}x^+}(p^{\hat{i}} + \text{f}p^+ \partial_{p^{\hat{i}}}), \quad (3.20)$$

$$J^{\hat{i}\hat{j}} = l^{\hat{i}\hat{j}} + M^{\hat{i}\hat{j}}, \quad l^{\hat{i}\hat{j}} \equiv p^{\hat{i}} \partial_{p^{\hat{j}}} - p^{\hat{j}} \partial_{p^{\hat{i}}}, \quad M^{\hat{i}\hat{j}} \equiv \theta^Z \bar{\gamma}^{-\hat{i}\hat{j}} \theta^{\bar{Z}}. \quad (3.21)$$

where we use notation $p^2 = p^{\hat{i}} p^{\hat{i}}$, $\partial_p^2 = \partial_{p^{\hat{i}}} \partial_{p^{\hat{i}}}$. These generators satisfy commutation relations of the isometry symmetries algebra of $4d$ plane wave background (see Appendix C).

Proper conformal generators:

$$D = -2\partial_{p^+} p^+ - \partial_{p^{\hat{i}}} p^{\hat{i}} + 2\theta^Z \bar{\gamma}^- \theta^{\bar{Z}} - 1, \quad (3.22)$$

$$C = e^{-2\text{if}x^+} \left(\frac{p^2}{2p^+} + \frac{\text{f}^2}{2} p^+ \partial_p^2 - \text{f} \partial_{p^{\hat{i}}} p^{\hat{i}} + \text{f} \right), \quad (3.23)$$

$$\begin{aligned} C^{\hat{i}} &= e^{-\text{if}x^+} \left(-\partial_{p^+} p^{\hat{i}} - \frac{1}{2p^+} \partial_{p^{\hat{i}}} p^2 + M^{\hat{i}\hat{j}} \frac{p^{\hat{j}}}{p^+} + (\theta^Z \bar{\gamma}^- \theta^{\bar{Z}} - 1) \frac{p^{\hat{i}}}{p^+} \right. \\ &\quad \left. + \text{f} \left(-\frac{1}{2} \partial_p^2 p^{\hat{i}} + \partial_{p^{\hat{i}}} (\partial_{p^+} p^+ + \partial_{p^{\hat{i}}} p^{\hat{i}} - \theta^Z \bar{\gamma}^- \theta^{\bar{Z}}) - M^{\hat{i}\hat{j}} \partial_{p^{\hat{j}}} \right) \right), \end{aligned} \quad (3.24)$$

$$\begin{aligned} K^- &= \frac{1}{4p^+} \partial_p^2 p^2 + (\partial_{p^+} + \frac{1}{p^+} (1 - \theta^Z \bar{\gamma}^- \theta^{\bar{Z}})) (\partial_{p^+} p^+ + \partial_{p^{\hat{i}}} p^{\hat{i}} - \theta^Z \bar{\gamma}^- \theta^{\bar{Z}}) \\ &\quad + \frac{1}{2p^+} M^{\hat{i}\hat{j}} l^{\hat{i}\hat{j}} + \frac{1}{2p^+} M^{\hat{i}\hat{j}} M^{\hat{i}\hat{j}}. \end{aligned} \quad (3.25)$$

Generators of the isometry symmetries and proper conformal symmetries form commutation relations of the $so(4, 2)$ algebra taken to be in the plane wave basis.

Generators of R-symmetries $so(6)$ algebra:

$$J^{i'j'} = \theta^Z \bar{\gamma}^{-i'j'} \theta^{\bar{Z}}, \quad J^{Z\bar{Z}} = \theta^Z \bar{\gamma}^- \theta^{\bar{Z}} - 1, \quad (3.26)$$

$$J^{Zi'} = \frac{p^+}{2} \theta^Z \bar{\gamma}^{-Zi'} \theta^{\bar{Z}}, \quad J^{\bar{Z}i'} = \frac{1}{2p^+} \theta^{\bar{Z}} \bar{\gamma}^{-\bar{Z}i'} \theta^{\bar{Z}}. \quad (3.27)$$

The hermitian properties of these generators are fixed to be $J^{i'j'\dagger} = -J^{j'i'}$, $J^{Zi'\dagger} = -J^{\bar{Z}i'}$, $J^{Z\bar{Z}\dagger} = J^{Z\bar{Z}}$.

Supercharges:

$$\Omega^+ = 2\bar{\gamma}^- (p^+ \theta^Z + \theta^{\bar{Z}}), \quad (3.28)$$

$$\Omega^- = 2T^{\hat{i}} \bar{\gamma}^{\hat{i}} (\theta^Z + \frac{1}{p^+} \theta^{\bar{Z}}), \quad \bar{\Omega}^- = 2\bar{T}^{\hat{i}} \bar{\gamma}^{\hat{i}} (\theta^Z + \frac{1}{p^+} \theta^{\bar{Z}}), \quad (3.29)$$

$$S^- = 2(D - 1)(\theta^Z + \frac{1}{p^+} \theta^{\bar{Z}}) + (l^{\hat{i}\hat{j}} + 2M^{\hat{i}\hat{j}}) \gamma^{\hat{i}\hat{j}} (\theta^Z + \frac{1}{p^+} \theta^{\bar{Z}}). \quad (3.30)$$

The hermitian properties of the supercharges are fixed to be $\Omega^{+\dagger} = \Omega^+$, $\Omega^{-\dagger} = \bar{\Omega}^-$, $S^{-\dagger} = -S^-$.

Transformations of fields defined by $\delta_G \phi = [\phi, G_{f.t.}]|_{eq. x^+}$ take then the form

$$\delta_G \phi = G \phi. \quad (3.31)$$

4 Oscillator realization of superconformal symmetries

The expressions for generators above-given are defined in terms of differential operators acting on wave function taken to be in momentum space. The representation for generators can be simplified by passing to the oscillators representation of wave function.

Let $\phi(p, p^+)$ be square integrable solution to free equations of motion for fields of plane wave $\mathcal{N} = 4$ SYM corresponding to the $p^+ > 0$, (dependence on $\theta^{\bar{Z}}$ is implicit)

$$i\partial_{x^+}\phi = -P^-\phi, \quad p^+ > 0. \quad (4.32)$$

Transformation of this wave function from momentum space to the oscillator basis wave function denoted by $|\phi(a, p^+)\rangle$ is fixed to be

$$|\phi(a, p^+)\rangle = \int d^2p \, e^F \phi(p, p^+) |0\rangle, \quad F \equiv -\frac{1}{2f p^+} p^{\hat{i}} p^{\hat{i}} + \sqrt{\frac{2}{f p^+}} a^{\hat{i}} p^{\hat{i}} - \frac{1}{2} a^{\hat{i}} a^{\hat{i}}, \quad (4.33)$$

$f > 0$, where $a^{\hat{i}}$ is creation operator and $|0\rangle$ is the Fock space vacuum defined by $\bar{a}^{\hat{i}}|0\rangle = 0$, $[\bar{a}^{\hat{i}}, a^{\hat{j}}] = \delta^{\hat{i}\hat{j}}$. These formulas imply the following realization of operators $a^{\hat{i}}$ and $\bar{a}^{\hat{i}}$

$$a^{\hat{i}} = \frac{1}{\sqrt{2f p^+}} (p^{\hat{i}} - f p^+ \partial_{p^{\hat{i}}}), \quad \bar{a}^{\hat{i}} = \frac{1}{\sqrt{2f p^+}} (p^{\hat{i}} + f p^+ \partial_{p^{\hat{i}}}). \quad (4.34)$$

Making use of these relation it is easy to see that the solution to the equation (4.32) can be presented in the following elegant form

$$|\phi(a, p^+)\rangle = \sum_{n=0}^{\infty} e^{-if(n+1)x^+} a^{\hat{i}_1} \dots a^{\hat{i}_n} \phi^{\hat{i}_1 \dots \hat{i}_n}(p^+) |0\rangle. \quad (4.35)$$

Taking into account a realization of the differential operator ∂_{p^+} on the oscillator wave function $|\phi(a, p^+)\rangle$

$$\left(\partial_{p^+} + \frac{1}{4p^+} (a^{\hat{i}} a^{\hat{i}} - \bar{a}^{\hat{i}} \bar{a}^{\hat{i}} - 2)\right) |\phi(a, p^+)\rangle = \int d^2p \, e^F \partial_{p^+} \phi(p, p^+) \quad (4.36)$$

and the formulas (4.34) it is easy to transform the generators to the oscillator basis. For the generators of the isometry symmetries we get the representation

$$P^+ = p^+, \quad P^- = -f(a^{\hat{i}} \bar{a}^{\hat{i}} + 1), \quad (4.37)$$

$$T^{\hat{i}} = \sqrt{2f p^+} e^{-ifx^+} a^{\hat{i}}, \quad \bar{T}^{\hat{i}} = \sqrt{2f p^+} e^{ifx^+} \bar{a}^{\hat{i}}, \quad (4.38)$$

$$J^{\hat{i}\hat{j}} = a^{\hat{i}} \bar{a}^{\hat{j}} - a^{\hat{j}} \bar{a}^{\hat{i}} + \theta^{\bar{Z}} \bar{\gamma}^{-\hat{i}\hat{j}} \theta^{\bar{Z}}. \quad (4.39)$$

Oscillator representation for generators of the proper conformal transformations takes the form

$$D = -2\partial_{p^+} p^+ + 2\theta^{\bar{Z}} \bar{\gamma}^{-\hat{i}\hat{i}} \theta^{\bar{Z}} - 1, \quad (4.40)$$

$$C = f e^{-2ifx^+} a^{\hat{i}} a^{\hat{i}}, \quad (4.41)$$

$$C^{\hat{i}} = -\partial_{p^+} T^{\hat{i}} - \frac{1}{2\mathfrak{f}p^+} (\bar{T}^{\hat{i}} C + T^{\hat{i}} P^-) + \frac{1}{p^+} \theta^Z \bar{\gamma}^- \gamma^{\hat{i}} \bar{\gamma}^{\hat{j}} \theta^{\bar{Z}} T^{\hat{j}}, \quad (4.42)$$

$$\begin{aligned} K^- = & \frac{1}{8\mathfrak{f}^2 p^+} \{C, \bar{C}\} - \frac{1}{4\mathfrak{f}^2 p^+} (P^-)^2 + (\partial_{p^+} - \frac{1}{p^+} \theta^Z \bar{\gamma}^- \theta^{\bar{Z}}) (\partial_{p^+} p^+ + 1 - \theta^Z \bar{\gamma}^- \theta^{\bar{Z}}) \\ & + \frac{1}{p^+} M^{\hat{i}\hat{j}} a^{\hat{i}} \bar{a}^{\hat{j}} + \frac{1}{2p^+} M^{\hat{i}\hat{j}} M^{\hat{i}\hat{j}} + \frac{3}{4p^+}. \end{aligned} \quad (4.43)$$

The oscillator representation for the supercharges is obtainable by inserting the above-given expressions for bosonic generators into formulas (3.28)-(3.30). Because of certain attractive features it seems that the oscillators representation is fruitful direction to go. Interesting discussion of application of the oscillator construction to the plane wave physics may be found in [28]⁵.

Acknowledgments. This work was supported by the INTAS project 991590, by the RFBR Grant 02-02-17067 and RFBR Grant for Leading Scientific Schools 00-15-96566.

Appendix A: Lagrangian of plane wave $\mathcal{N} = 4$ SYM in terms of component fields on space-time. Light cone gauge fixed Lagrangian can be presented as follows

$$\mathcal{L} = \mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4, \quad (4.44)$$

where $\mathcal{L}_2$ describes quadratic part of Lagrangian while $\mathcal{L}_3$ and $\mathcal{L}_4$ describe 3-point and 4-point interaction vertices respectively. The quadratic part $\mathcal{L}_2$ is given by

$$\mathcal{L}_2 = \text{Tr} \left[\frac{1}{2} A^I \square A^I - \frac{\mathfrak{i}}{4} \psi^{\oplus} \frac{\bar{\gamma}^+}{\partial^+} \square \psi^{\oplus} \right], \quad (4.45)$$

where the standard covariant D'Alembertian operator in 4d plane wave background

$$ds^2 = 2dx^+ dx^- - \mathfrak{f}^2 x^{\hat{i}} x^{\hat{i}} dx^+ dx^+ + dx^{\hat{i}} dx^{\hat{i}}, \quad \hat{i} = 1, 2, \quad (4.46)$$

is given by

$$\square = 2\partial_{x^+} \partial_{x^-} + \partial_{x^{\hat{i}}} \partial_{x^{\hat{i}}} + \mathfrak{f}^2 x^{\hat{i}} x^{\hat{i}} \partial_{x^-} \partial_{x^-}. \quad (4.47)$$

The 3-point and 4-point interaction vertices are given by

$$\mathcal{L}_3 = \text{Tr} \left[-[A^I, A^J] \partial^I A^J - \partial^J A^J \frac{1}{\partial^+} [A^I, \partial^+ A^I] \right] \quad (4.48)$$

$$+ \mathfrak{i} \partial^I A^I \frac{1}{\partial^+} (\psi^{\oplus} \bar{\gamma}^+ \psi^{\oplus}) + \frac{\mathfrak{i}}{4} [A^I, \psi^{\oplus}] \frac{\bar{\gamma}^+ \gamma^I \bar{\gamma}^J}{\partial^+} \partial^J \psi^{\oplus} + \frac{\mathfrak{i}}{4} \partial^J \psi^{\oplus} \frac{\bar{\gamma}^+ \gamma^J \bar{\gamma}^I}{\partial^+} [A^I, \psi^{\oplus}],$$

$$\begin{aligned} \mathcal{L}_4 = & \text{Tr} \left[-\frac{1}{4} [A^I, A^J]^2 - \frac{1}{2} \left(\frac{1}{\partial^+} [A^J, \partial^+ A^J] \right)^2 \right. \\ & + \frac{\mathfrak{i}}{\partial^+} (\psi^{\oplus} \bar{\gamma}^+ \psi^{\oplus}) \frac{1}{\partial^+} [A^I, \partial^+ A^I] + \frac{\mathfrak{i}}{4} [A^I, \psi^{\oplus}] \frac{\bar{\gamma}^+ \gamma^I \bar{\gamma}^J}{\partial^+} [A^J, \psi^{\oplus}] \\ & \left. + \frac{1}{2} \left(\frac{1}{\partial^+} (\psi^{\oplus} \bar{\gamma}^+ \psi^{\oplus}) \right)^2 \right], \quad \partial^I = \partial_{x^I}, \quad \partial^+ = \partial_{x^-}, \quad \partial^- = \partial_{x^+}. \end{aligned} \quad (4.49)$$

⁵Realization of conformal and Heisenberg algebras in plane wave CFT correspondence may be found in [29].

The physical fields A^I , $\psi^\oplus$ being Lie algebra valued in the adjoint representation of gauge group are assumed to be anti-hermitian $(A^I)^\dagger = -A^I$, $\psi^{\oplus\dagger} = -\psi^\oplus$. The Tr denotes minus trace $\text{Tr}Y \equiv -\text{Tr}Y$. All fields are assumed to be independent of six coordinates x^M : $\partial_{x^M}A^I = 0$, $\partial_{x^M}\psi^\oplus = 0$.

Appendix B: Component form of the action of $N = 4$ SYM in terms of momentum modes. Because a formulation of action in terms of momentum modes of fields is more convenient in superfield approach we make the Fourier transformation

$$\phi(x) = i \int \frac{d^3p}{(2\pi)^{3/2}} e^{i(p^+x^- + x^i p^i)} \phi(x^+, p), \quad d^3p \equiv dp^1 dp^2 dp^+, \quad (4.50)$$

so that the Fourier modes satisfy the hermitian conjugation rules

$$A^I(x^+, p)^\dagger = A^I(x^+, -p), \quad \psi^\oplus(x^+, p)^\dagger = \psi^\oplus(x^+, -p). \quad (4.51)$$

In terms of momentum modes the quadratic and the cubic parts of the action (4.45), (4.48) take the form

$$S_2 = \text{Tr} \int dx^+ d^3p A^I(-p) p^+ (i\partial^- + P^-) A^I(p) - \frac{i}{2} \psi^\oplus(-p) \bar{\gamma}^+ (i\partial^- + P^-) \psi^\oplus(p), \quad (4.52)$$

$$S_3 = \text{Tr} \int d\Gamma_3(p) \left(\frac{2}{p_3^+} \mathbf{P}^K \delta^{IJ} A^I(p_1) A^J(p_2) A^K(p_3) \right. \\ \left. - \frac{p_1^+ - p_2^+}{2p_1^+ p_2^+ p_3^+} \mathbf{P}^I \psi(p_1) \bar{\gamma}^+ \psi(p_2) A^I(p_3) - \frac{1}{2p_1^+ p_2^+} \mathbf{P}^I \psi(p_1) \bar{\gamma}^{+IJ} \psi(p_2) A^J(p_3) \right), \quad (4.53)$$

where the momentum $\mathbf{P}^I$ is defined in (2.12), while the measure $d\Gamma_3(p)$ is given by

$$d\Gamma_3(p) = \frac{1}{(2\pi)^{3/2}} \delta^{(3)}\left(\sum_{a=1}^3 p_a\right) \prod_{a=1}^3 d^3p_a. \quad (4.54)$$

In these formulas we keep dependence on six coordinates x^M . To get expressions corresponding to 4d SYM one needs to apply rule $\mathbf{P}^M = 0$.

Appendix C: Commutation relations of the $psu(2, 2|4)$ superalgebra in plane wave basis. These commutation relations for anti-hermitian bosonic generators were worked out in [22]. In this paper we use the basis in which some of the generators have different hermitian properties. This is to say that $P^\pm$, K^- are taken to be hermitian, while the hermitian properties of the remaining generators are fixed to be $T^{\hat{i}\dagger} = \bar{T}^{\hat{i}}$, $J^{\hat{i}\hat{j}\dagger} = -J^{\hat{i}\hat{j}}$, $D^\dagger = -D$, $C^\dagger = \bar{C}$, $C^{\hat{i}\dagger} = -C^{\hat{i}6}$.

In plane wave basis all generators, by definition, are eigenvectors of P^- and D . Therefore given generator G we introduce $q_G(P^-)$ and $q_G(D)$ charges which are defined by commutators

$$[P^-, G] = -f q_G(P^-) G, \quad [D, G] = q_G(D) G. \quad (4.55)$$

These charges for generators of the $psu(2, 2|4)$ superalgebra are given in Table.

⁶Denoting by G_{ah} the bosonic anti-hermitian generators used in [22] we have the following relations: $P^\pm = -iP_{ah}^\pm$, $T^{\hat{i}} = -iT_{ah}^{\hat{i}}$, $\bar{T}^{\hat{i}} = -i\bar{T}_{ah}^{\hat{i}}$, $J^{\hat{i}\hat{j}} = J_{ah}^{\hat{i}\hat{j}}$, $D = D_{ah}$, $C = iC_{ah}$, $\bar{C} = i\bar{C}_{ah}$, $C^{\hat{i}} = C_{ah}^{\hat{i}}$, $\bar{C}^{\hat{i}} = \bar{C}_{ah}^{\hat{i}}$, $K^- = iK_{ah}^-$. Here we use the same supercharges $\Omega^\pm$ as in [22], while S^- is multiplied by factor i .

Generators	P^+	$T^{\hat{i}}$	$\bar{T}^{\hat{i}}$	K^-	$C^{\hat{i}}$	$\bar{C}^{\hat{i}}$	C	$\bar{C}$	Ω^-	$\bar{\Omega}^-$	Ω^+	S^-
q - charges												
$q_G(P^-)$ charge	0	1	-1	0	1	-1	2	-2	1	-1	0	0
$q_G(D)$ charge	-2	-1	-1	2	1	1	0	0	0	0	-1	1

Note that the q -charges of the generators $J^{\hat{i}\hat{j}}$ and J^{MN} are equal to zero. Remaining (anti)commutators can be collected in several groups.

Commutators between elements of isometry algebra are

$$[\bar{T}^{\hat{i}}, T^{\hat{j}}] = 2f\delta^{\hat{i}\hat{j}}P^+, \quad [T^{\hat{i}}, J^{\hat{j}\hat{k}}] = \delta^{\hat{i}\hat{j}}T^{\hat{k}} - \delta^{\hat{i}\hat{k}}T^{\hat{j}}. \quad (4.56)$$

Commutators between proper conformal generators are given by

$$[\bar{C}^{\hat{i}}, C^{\hat{j}}] = 2\delta^{\hat{i}\hat{j}}K^-, \quad [\bar{C}, C] = -4fP^-, \quad [\bar{C}^{\hat{i}}, C] = 2fC^{\hat{i}}. \quad (4.57)$$

Commutators between generators of isometry algebra and proper conformal generators take the form

$$[P^+, K^-] = D, \quad [P^+, C^{\hat{i}}] = T^{\hat{i}}, \quad [T^{\hat{i}}, K^-] = C^{\hat{i}}, \quad (4.58)$$

$$[T^{\hat{i}}, C^{\hat{j}}] = \delta^{\hat{i}\hat{j}}C, \quad [T^{\hat{i}}, \bar{C}] = -2f\bar{T}^{\hat{i}}, \quad (4.59)$$

$$[\bar{C}^{\hat{i}}, T^{\hat{j}}] = \delta^{\hat{i}\hat{j}}(P^- + fD) + 2fJ^{\hat{i}\hat{j}}. \quad (4.60)$$

Commutators between isometry algebra and supercharges are

$$[T^{\hat{i}}, \bar{\Omega}^-] = f\bar{\gamma}^{+\hat{i}}\Omega^+, \quad [T^{\hat{i}}, S^-] = \gamma^{\hat{i}}\Omega^-, \quad [P^+, S^-] = \gamma^+\Omega^+, \quad (4.61)$$

$$[\Omega^\pm, J^{\hat{i}\hat{j}}] = \frac{1}{2}\bar{\gamma}^{\hat{i}\hat{j}}\Omega^\pm, \quad [S^-, J^{\hat{i}\hat{j}}] = \frac{1}{2}\gamma^{\hat{i}\hat{j}}S^-.$$

Commutators between proper conformal generators and supercharges are

$$[K^-, \Omega^+] = -\frac{1}{2}\bar{\gamma}^-S^-, \quad [C^{\hat{i}}, \Omega^+] = -\frac{1}{2}\bar{\gamma}^{-\hat{i}}\Omega^-, \quad [C^{\hat{i}}, \bar{\Omega}^-] = -f\bar{\gamma}^{\hat{i}}S^-, \quad (4.62)$$

$$[C, \bar{\Omega}^-] = -2f\Omega^-. \quad (4.63)$$

Commutators of $so(6)$ algebra and commutators between generators of the $so(6)$ and supercharges in plane wave basis take the standard form.

Anticommutators between supercharges are

$$\{\Omega^+, \Omega^+\} = 2\bar{\gamma}^-P^+, \quad \{\Omega^-, \Omega^-\} = -2\bar{\gamma}^+C, \quad \{S^-, S^-\} = 4\gamma^+K^-, \quad (4.64)$$

$$\{\bar{\Omega}^-, \Omega^-\} = 2\bar{\gamma}^+P^- + f\bar{\gamma}^{+\hat{i}\hat{j}}J^{\hat{i}\hat{j}} - f\bar{\gamma}^{+MN}J^{MN}, \quad (4.65)$$

$$\{\Omega^-, \Omega^+\} = \bar{\gamma}^+\gamma^-\bar{\gamma}^{\hat{i}}T^{\hat{i}}, \quad \{\Omega^-, S^-\} = -2\bar{\gamma}^{+\hat{i}}C^{\hat{i}}, \quad (4.66)$$

$$\{S^-, \Omega^+\} = \gamma^+\bar{\gamma}^-D + \frac{1}{2}\gamma^+\bar{\gamma}^-\gamma^{\hat{i}\hat{j}}J^{\hat{i}\hat{j}} - \frac{1}{2}\gamma^+\bar{\gamma}^-\gamma^{MN}J^{MN}. \quad (4.67)$$

Modulo (anti)commutators obtainable from the ones above-given by applying hermitian conjugation, the remaining (anti)commutators are equal to zero.

Appendix D: Fierz identities. Often we use the following identities for γ -matrices⁷

$$\gamma_{\alpha(\beta}^{\mu}\gamma_{\delta\gamma)}^{\mu} = 0, \quad \delta_{\alpha'}^{\alpha}\delta_{\beta'}^{\beta} - \delta_{\beta'}^{\alpha}\delta_{\alpha'}^{\beta} = \frac{1}{48}(\gamma^{\mu\nu\rho})^{\alpha\beta}(\gamma^{\mu\nu\rho})_{\alpha'\beta'}, \quad (4.68)$$

$$\delta_{\beta}^{\alpha}\delta_{\gamma}^{\delta} - \frac{1}{2}(\gamma^{mn})^{\alpha}_{\beta}(\gamma^{mn})^{\delta}_{\gamma} + \frac{1}{2}(\gamma^{MN})^{\alpha}_{\beta}(\gamma^{MN})^{\delta}_{\gamma} + (\beta \leftrightarrow \gamma) = 2\gamma_{\beta\gamma}^m(\gamma^m)^{\alpha\delta}. \quad (4.69)$$

The identities (4.68) are well known, while the identity (4.69) is derivable from the 1st identity in (4.68). From the 2nd identity (4.68) we can get the following ‘generating’ identity

$$8\psi\bar{\gamma}^{-}\theta^Z\theta^Z\bar{\gamma}^{-}\varphi = \psi\bar{\gamma}^{-\bar{Z}\hat{I}}\varphi\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z. \quad (4.70)$$

Inserting in (4.70) $\psi = \theta^Z\gamma^{Z\hat{I}}$, $\varphi = \gamma^{Z\hat{J}}\theta^Z$ and $\psi = \theta^Z\gamma^{Zi'}$, $\varphi = \gamma^{+Z}\psi^{\oplus Z}$ we get the following respective identities

$$\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z\theta^Z\bar{\gamma}^{-Z\hat{J}}\theta^Z = \frac{1}{6}\delta^{\hat{I}\hat{J}}\theta^Z\bar{\gamma}^{-Z\hat{K}}\theta^Z\theta^Z\bar{\gamma}^{-Z\hat{K}}\theta^Z, \quad (4.71)$$

$$\theta^Z\bar{\gamma}^{-Zi'}\theta^Z\theta^Z\bar{\gamma}^Z\psi^{\oplus Z} = -\frac{1}{6}\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z\theta^Z\bar{\gamma}^{\hat{I}Zi'}\psi^{\oplus Z}. \quad (4.72)$$

Multiplying the identity (4.69) by γ^{+} , γ^{-} and then by θ^Z , θ^Z gives the identity

$$(\bar{\gamma}^{\hat{I}}\theta^Z)_{\alpha}(\bar{\gamma}^{\hat{I}}\theta^Z)_{\beta} = -\frac{1}{8}(\bar{\gamma}^{+\bar{Z}\hat{I}})_{\alpha\beta}\theta^Z\bar{\gamma}^{-Zi'}\theta^Z. \quad (4.73)$$

Appendix E: Integrals over anti-commuting variables. The measure $d^4\theta^Z$ is normalized to be

$$\int d^4\theta^Z \theta^{Z\alpha}\theta^{Z\beta}\theta^{Z\gamma}\theta^{Z\delta} = \frac{1}{64}(\gamma^{+\bar{Z}\hat{I}})^{\alpha\beta}(\gamma^{+\bar{Z}\hat{I}})^{\gamma\delta}. \quad (4.74)$$

Note that the l.h.s. of this formula is antisymmetric in spinor indices $\alpha, \beta, \gamma, \delta$. Making use of the 1st identity (4.68) one can make sure that the r.h.s is also antisymmetric in $\alpha, \beta, \gamma, \delta$. The fact that the integral (4.74) can be cast into the form above-given can be proved by using identity

$$\theta^{Z\alpha}\theta^{Z\beta} = \frac{1}{16}(\gamma^{+\bar{Z}\hat{I}})^{\alpha\beta}\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z, \quad (4.75)$$

and identity (4.71). All remaining integrals are derivable from the basic integral (4.74) in a straightforward way:

$$\int d^4\theta^Z (\theta^Z\bar{\gamma}^{-}\theta^{Z\dagger})^4 = \frac{1}{16}(\theta^{Z\dagger}\bar{\gamma}^{-\bar{Z}\hat{I}}\theta^{Z\dagger})^2, \quad (4.76)$$

$$\int d^4\theta^Z \theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z \theta^Z\bar{\gamma}^{-Z\hat{J}}\theta^Z = 4\delta^{\hat{I}\hat{J}}, \quad (4.77)$$

$$\int d^4\theta^Z \theta^{Z\alpha}\theta^{Z\beta} \theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z = \frac{1}{4}(\gamma^{+\bar{Z}\hat{I}})^{\alpha\beta}. \quad (4.78)$$

Taking these relations into account the expression for δ -function is fixed to be

$$\delta(\theta^Z) = \frac{1}{24}\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z\theta^Z\bar{\gamma}^{-Z\hat{I}}\theta^Z. \quad (4.79)$$

⁷ $m, n = 0, 1, 2, 9$ are $\text{so}(3,1)$ indices; $M, N = 3, \dots, 8$ are R-symmetry $\text{so}(6)$ indices; $\alpha, \beta, \gamma, \delta = 1, \dots, 16$ are spinor indices. We use convention for γ matrices following Ref. [22].

References

- [1] M. Blau, J. Figueroa-O’Farrill, C. Hull and G. Papadopoulos, JHEP **0201**, 047 (2002) [hep-th/0110242]; Class. Quant. Grav. **19**, L87 (2002); hep-th/0201081.
- [2] R. R. Metsaev, Nucl. Phys. B **625**, 70 (2002) [hep-th/0112044].
- [3] D.Berenstein, J.M.Maldacena, H.Nastase, JHEP**0204**, 013(2002) hep-th/0202021
- [4] R. R. Metsaev and A. A. Tseytlin, Phys. Rev. D **65**, 126004 (2002) [hep-th/0202109].
- [5] C.S. Chu, V.V. Khoze, G. Travaglini, JHEP **0209**, 054 (2002) hep-th/0206167
- [6] N.Beisert, C.Kristjansen, J.Plefka, G.W.Semenoff, M.Staudacher, hep-th/0208178
- [7] N. R. Constable, D. Z. Freedman, M. Headrick and S. Minwalla, hep-th/0209002.
- [8] N. Beisert, “BMN operators and superconformal symmetry,” arXiv:hep-th/0211032.
- [9] C. Kristjansen, J. Plefka, G. W. Semenoff and M. Staudacher, Nucl. Phys. B **643**, 3 (2002) [hep-th/0205033].
- [10] M. Bianchi, B. Eden, G. Rossi and Y. S. Stanev, Nucl. Phys. B **646**, 69 (2002) [arXiv:hep-th/0205321].
- [11] A. Parnachev and A. V. Ryzhov, JHEP **0210**, 066 (2002) [arXiv:hep-th/0208010].
- [12] J. A. Minahan and K. Zarembo, arXiv:hep-th/0212208.
- [13] C. S. Chu and P. M. Ho, Nucl. Phys. B **636**, 141 (2002) [hep-th/0203186].
- [14] K. Skenderis and M. Taylor, “Branes in AdS and pp-wave spacetimes,” JHEP **0206**, 025 (2002) [hep-th/0204054]; arXiv:hep-th/0211011; arXiv:hep-th/0212184.
- [15] M. Billo and I. Pesando, Phys. Lett. B **536**, 121 (2002) [hep-th/0203028].
A. Dabholkar and S. Parvizi, Nucl. Phys. B **641**, 223 (2002) [hep-th/0203231].
P.Bain, K.Peeters and M.Zamaklar, hep-th/0208038. Y.Michishita, hep-th/0206131.
- [16] D. S. Bak, hep-th/0204033.
- [17] O. Bergman, M. R. Gaberdiel and M. B. Green, hep-th/0205183.
- [18] S. Corley and S. Ramgoolam, arXiv:hep-th/0212166.
- [19] M. Blau, JHEP **0011**, 023 (2000) [hep-th/0005098].
- [20] K. Okuyama, JHEP **0211**, 043 (2002) [arXiv:hep-th/0207067].
- [21] I.Y.Park, A.Sadrzadeh and T.A.Tran, Phys.Lett.B **497**, 303 (2001) hep-th/0010116.
- [22] R. R. Metsaev, arXiv:hep-th/0211178.
- [23] S. R. Das, C. Gomez and S. J. Rey, Phys. Rev. D **66**, 046002 (2002) hep-th/0203164.
E. Kiritsis and B. Pioline, JHEP **0208**, 048 (2002) [hep-th/0204004].
R.G.Leigh, K.Okuyama and M.Rozali, Phys.Rev.D**66**,046004(2002) hep-th/0204026.
D. Berenstein and H. Nastase, hep-th/0205048.
G. Siopsis, Phys. Lett. B **545**, 169 (2002) [hep-th/0205302]; arXiv:hep-th/0212165.
S.Dobashi, H.Shimada, T.Yoneya, hep-th/0209251; V.Sahakian, hep-th/0209179.
- [24] A. Santambrogio and D. Zanon, Phys. Lett. B **545**, 425 (2002) hep-th/0206079.
- [25] L. Brink, M. B. Green and J. H. Schwarz, Nucl. Phys. B **223**, 125 (1983).
- [26] L. Brink, O. Lindgren and B. E. Nilsson, Nucl. Phys. B **212**, 401 (1983).
S. Mandelstam, Nucl. Phys. B **213**, 149 (1983).
- [27] M. B. Green and J. H. Schwarz, Nucl. Phys. B **243**, 475 (1984).
- [28] S. Fernando, M. Gunaydin and O. Pavlyk, JHEP **0210**, 007 (2002) hep-th/0207175
- [29] S. R. Das and C. Gomez, JHEP **0207**, 016 (2002) [hep-th/0206062].
G. Arutyunov and E. Sokatchev, JHEP **0208**, 014 (2002) [hep-th/0205270].